

Virus 연구 및 백신 개발 with WES

Residual bovine serum albumin (BSA) quantitation in vaccines using automated Capillary Western technology

Author links open overlay panel [John W. Loughney](#) [Catherine Lancaster](#) [ShaHaRichard R. Rustandi](#)
Analytical Biochemistry [Volume 461](#), 15 September 2014, Pages 49-56

[More Information >>](#)

Inactivated Rabies Virus-Vectored Immunocontraceptive Vaccine in a Thermo-Responsive Hydrogel Induces High and Persistent Antibodies against Rabies, but Insufficient Antibodies against Gonadotropin-Releasing Hormone for Contraception

by [Xianfu Wu](#)^{1,*}, [Yong Yang](#)^{1,2}, [Chantal Kling](#)^{1,3}, [Laurie Seigler](#)^{1,2}, [Nadia F. Gallardo-Romero](#)¹, [Brock E. Martin](#)¹, [Todd G. Smith](#)¹ and [Victoria A. Olson](#)¹

Vaccines 2019, 7(3), 73;

[More Information >>](#)

SKP2 attenuates autophagy through Beclin1-ubiquitination and its inhibition reduces MERS-Coronavirus infection

Nils C. Gassen, Daniela Niemeyer, Doreen Muth, Victor M. Corman, Silvia Martinelli, Alwine Gassen, Kathrin Hafner, Jan Papias, Kirstin Mösbauer, Andreas Zellner, Anthony S. Zannas, Alexander Herrmann, Florian Holsboer, Ruth Brack-Werner, Michael Boshart, Bertram Müller-Myhsok, Christian Drosten, Marcel A. Müller & Theo Rein

Nature Communications volume 10, Article number: 5770 (2019) Cite this article

[More Information >>](#)

Automated Western blot, Wes



High Sensitivity: 일반 WB보다 8배 높은 sensitivity

Less Sample & less Ab: 0.6ug Sample
(최소농도 0.25mg/ml)

Multiplex Western: 최대 4개 targets 동시 분석 / sample

3hrs Run time: from loading to result

Full Automation: 높은 재현성

분석서비스 및 데모 가능

